arXiv:1504.05007v1 [math.FA] 20 Apr 2015

Integral triangular operators and Friedrichs model

Lev Sakhnovich

99 Cove ave., Milford, CT, 06461, USA.
E-mail: lsakhnovich@gmail.com

Abstract

In the present paper we investigate a semi-group of triangular integral operators V_β, which is an analogue of the semi-group of the fractional integral operators J^β. With the help of these semi-groups, we construct and study two classes of triangular Friedrichs models A_β and B_β, respectively. Using generalized wave operators we prove that A_β and B_β are linearly similar to a self-adjoint operator with absolutely continuous spectrum.

Mathematical Subject Classification (2010): 47A45; 47A15, 47A46.

Keywords: Triangular representation, semi-group, Friedrichs model, generalized wave operators.

1 Introduction

1. In the present paper we investigate the triangular integral operators

$$V_\beta f = \frac{1}{\Gamma(\beta)} \int_0^x E_\beta(x-t) f(t)\,\mathrm{d}t, \quad \Re\beta > 0, \quad f(x) \in L^2(0,\omega), \tag{1.1}$$

where $\Gamma(z)$ is Euler gamma function,

$$E_\beta(x) = \int_0^\infty \frac{1}{\Gamma(s)} \mathrm{e}^{-Cs} s^{\beta-1} x^{s-1}\,\mathrm{d}s, \quad C \in \mathbb{C}, \tag{1.2}$$

and $\mathbb{C}$ stands for the complex plane. We shall show that the operators V_β form a semi-group, which is an analogue of the semi-group of the fractional integral operators

$$J^\beta f = \frac{1}{\Gamma(\beta)} \int_0^x (x-t)^{\beta-1} f(t)\,\mathrm{d}t, \quad \Re\beta > 0, \quad f(x)\in L^2(0,\omega). \tag{1.3}$$

With the help of the semi-groups of operators J^β and V_β we construct and study two classes of triangular Friedrichs models A_β and B_β, respectively. Using generalized wave operators we prove that A_β and B_β are linearly similar to a self-adjoint operator with absolutely continuous spectrum.

Lemma 1.1 *The function $E_\beta(x)$ is continuous in the domain $(0,\omega]$ and*

$$E_\beta(x) = \frac{\Gamma(\beta+1)}{x|\ln(x)|^{\beta+1}} \left(1 + O\left(\frac{1}{\ln(x)}\right)\right), \quad x\to 0. \tag{1.4}$$

Proof. It follows from (1.2) that the function $E_\beta(x)$ is continuous in the domain $(0,\omega]$ and

$$E_\beta(x) = \int_0^1 \frac{1}{\Gamma(s)} \mathrm{e}^{-Cs} s^{\beta-1} x^{s-1}\,\mathrm{d}s + O(1). \tag{1.5}$$

Using relations

$$\frac{1}{\Gamma(s)} = s + O(s^2), \quad \mathrm{e}^{-Cs} = 1 + O(s) \qquad (s\to 0), \tag{1.6}$$

we obtain

$$E_\beta(x) = \frac{1}{x}\int_0^1 s^\beta\big(1 + O(s)\big)\mathrm{e}^{-s|\ln(x)|}\,\mathrm{d}s + O(1). \tag{1.7}$$

Equality (1.7) implies the equality (1.4). The lemma is proved. □

Proposition 1.2 *The operators V_β, defined by formula* (1.1), *are bounded in all spaces $L^p(0,\omega)$, $p\geq 1$.*

Proof. Indeed, according to (1.4) we have (see [9, p. 24])

$$\big\|V_\beta\big\|_p \leq m(\beta) = \int_0^\omega \big|E_\beta(x)/\Gamma(\beta)\big|\,\mathrm{d}x < \infty. \tag{1.8}$$

This proves the proposition. □

The operators V_β have a following important property [10].

Theorem 1.3 *The operators* V_β $(\Re\beta > 0)$ *defined by formula* (1.1), *form a semi-group, that is,*

$$V_\alpha V_\beta = V_{\alpha+\beta}, \quad \Re\alpha > 0, \quad \Re\beta > 0. \tag{1.9}$$

2. We introduce the following integro-differential operator

$$Rf = -\int_0^x f'(t)\ln(x-t)\,\mathrm{d}t, \quad f(x)\in L(0,\omega). \tag{1.10}$$

In the book [9, p. 73] we proved that the operator V_1 is a right inverse of R.

Proposition 1.4 *If* $\beta = 1$ *and* $C = -\Gamma'(1)$ *then*

$$RV_1\varphi = \varphi, \quad \varphi'(x)\in L(0,\omega). \tag{1.11}$$

3. By H_α we denote the space of all function such that $f(x)\in L^2(0,\omega)$ and $f(x) = 0$ when $x\in[0,\alpha]$. It is easy to see that the spaces H_α, $0 < \alpha\leq\omega$, are invariant subspaces of the operator V_β.

Definition 1.5 *A bounded operator* T *is unicellular if its lattice of invariant subspaces is totally ordered by inclusion.*

Theorem 1.6 *The operator* V_β, $0 < \beta\leq 1$, *defined by formula* (1.1) *is unicellular.*

Proof. Let us consider the inner product

$$\left(V_\beta^n f, g\right) = \int_0^\infty \frac{1}{\Gamma(s)}\mathrm{e}^{-Cs}s^{n\beta-1}\left(\int_0^\omega \int_0^x (x-t)^{s-1}f(t)\,\mathrm{d}t\,\overline{g(x)}\,\mathrm{d}x\right)\mathrm{d}s. \tag{1.12}$$

Assuming that $\left(V_\beta^n f, g\right) = 0$ we use the set of functions $\psi_n(x) = \mathrm{e}^{-x^2/2}H_{2n}(x)$, where $H_{2n}(x)$ are Hermite polynomials. The set of functions $\psi_n(x)$ is complete in the Hilbert space $L^2(0,\infty)$. We note that $H_{2n}(x)$ are even polynomials of the degree $2n$. Hence, the set of functions $\varphi_n(x) = \mathrm{e}^{-x^2/2}x^{2n}$ is complete in the Hilbert space $L^2(0,\infty)$. Changing the variable $s^{\beta/2} = u$ and taking into account the formulas (1.12) and $\left(V_\beta^n f, g\right) = 0$, we have

$$\int_0^\omega \int_0^x (x-t)^{s-1}f(t)\,\mathrm{d}t\,\overline{g(x)}\,\mathrm{d}x = 0, \quad 0 < \beta\leq 1. \tag{1.13}$$

Relation (1.13) can be written in the form

$$\int_0^\omega u^{s-1} \int_u^\omega f(x-u)\overline{g(x)}\,\mathrm{d}x\,\mathrm{d}u = 0. \tag{1.14}$$

It follows from (1.14) that

$$\int_u^\omega f(x-u)\overline{g(x)}\,\mathrm{d}x = 0. \tag{1.15}$$

By changing the variables $x = \omega - x_1$, $u = \omega - u_1$ we obtain

$$\int_0^{u_1} f(u_1 - x_1)\overline{g(\omega - x_1)}\,\mathrm{d}x_1 = 0. \tag{1.16}$$

Using well-known Titchmarsh's theorem ([11, Theorem 152]) we receive the following assertion: if $f(x) \in H_\alpha$ and $f(x) \notin H_\gamma$, when $\gamma > \alpha$, then the system $V_\beta^n f$ is complete in the space H_α. This proves the theorem. □

Remark 1.7 *Using Titchmarsh's theorem ([11, Theorem 152]) it is easy to prove that operator* J^β, $\beta > 0$, *is unicellular.*

2 Triangular integral operators with logarithmic type kernels

1. In the present section we shall consider the operators

$$S_\beta f = \beta \int_0^x f(t)\big|\ln(x-t)\big|^{-\beta-1} \frac{\mathrm{d}t}{x-t}, \quad \Re\beta > 0, \tag{2.1}$$

in the space $L^2(0,\omega)$, $0 < \omega < 1$. The operator S_β is the main term in the expression which defines the operator V_β (see (1.1), (1.2)). Thus, by investigating the operator S_β we also obtain some properties of the operator V_β. The operator S_β is of independent interest as well. Rewrite the operator S_β in the following form

$$S_\beta f = \frac{\mathrm{d}}{\mathrm{d}x} \int_0^x f(t)\big|\ln(x-t)\big|^{-\beta}\,\mathrm{d}t, \quad \Re\beta \geq 0. \tag{2.2}$$

We note that the operator S_β is well-defined by formula (2.2) in the case where $\Re\beta\geq 0$. Let us introduce the functions (see [8] and [10])

$$\widetilde{s}(\lambda) = \int_0^\omega \mathrm{e}^{\mathrm{i}t\lambda}\big|\ln(t)\big|^{-\beta}\,\mathrm{d}t, \quad \Re\beta\geq 0, \tag{2.3}$$

$$\widetilde{s}_1(\lambda) = \int_0^\omega \mathrm{e}^{\mathrm{i}t\lambda}\big|\ln(t)\big|^{-\beta}\big(1 - t/\omega\big)\,\mathrm{d}t, \quad \Re\beta\geq 0. \tag{2.4}$$

Theorem 2.1 *If* $\Re\beta\geq 0$ *and* $0 < \omega < 1$*, then the following asymptotic relations are valid:*

$$-\mathrm{i}\lambda\widetilde{s}(\lambda) = \big(\ln|\lambda|\big)^{-\beta}\big(1 + o(1)\big) - \mathrm{e}^{\mathrm{i}\lambda\omega}\big(-\ln\omega\big)^{-\beta}, \quad \lambda\to\pm\infty, \tag{2.5}$$

$$-\mathrm{i}\lambda\widetilde{s_1}(\lambda) = \big(\ln|\lambda|\big)^{-\beta}\big(1 + o(1)\big), \quad \lambda\to\pm\infty. \tag{2.6}$$

Proof. According to the Cauchy theorem we have

$$\int_\gamma \mathrm{e}^{\mathrm{i}t\lambda}\big(-\ln(t)\big)^{-\beta}\,\mathrm{d}t = 0, \tag{2.7}$$

where the curve γ is depicted by Fig.1.



Figure 1: Contour of integration

From (2.3) and (2.7) we obtain the equality

$$\widetilde{s}(\lambda) = \int_0^{\mathrm{i}\omega} \mathrm{e}^{\mathrm{i}t\lambda}\big(-\ln t\big)^{-\beta}\,\mathrm{d}t - \int_0^{\pi/2} \mathrm{e}^{\mathrm{i}t\lambda}\big(-\ln(t)\big)^{-\beta}\mathrm{i}t\,\mathrm{d}\varphi, \quad t = \omega\mathrm{e}^{\mathrm{i}\varphi}. \tag{2.8}$$

Let us consider the integral

$$\int_0^{\mathrm{i}\omega} \mathrm{e}^{\mathrm{i}t\lambda}\bigl(-\ln t\bigr)^{-\beta}\,\mathrm{d}t = \int_0^{\lambda\omega} \mathrm{e}^{-u}\bigl(-\ln u+\ln\lambda-\mathrm{i}\pi/2\bigr)^{-\beta}\mathrm{i}\,\mathrm{d}u/\lambda, \quad \lambda>0. \tag{2.9}$$

When λ tends to infinity, equality (2.9) yields

$$\int_0^{\mathrm{i}\omega} \mathrm{e}^{\mathrm{i}t\lambda}\bigl(-\ln t\bigr)^{-\beta}\,\mathrm{d}t = \frac{\mathrm{i}}{\lambda}\bigl(\ln\lambda\bigr)^{-\beta}\bigl(1+o(1)\bigr), \quad \lambda\to+\infty. \tag{2.10}$$

Integrating by parts we estimate the integral

$$\int_0^{\pi/2} \mathrm{e}^{\mathrm{i}t\lambda}\bigl(-\ln(t)\bigr)^{-\beta}\mathrm{i}t\,\mathrm{d}\varphi = -\frac{1}{\mathrm{i}\lambda}\Bigl(\mathrm{e}^{\mathrm{i}\lambda\omega}\bigl(-\ln\omega\bigr)^{-\beta}+o(1)\Bigr),\ \lambda\to+\infty, \tag{2.11}$$

where $t=\omega\mathrm{e}^{\mathrm{i}\varphi}$, $\lambda\to+\infty$. Relations (2.8), (2.10) and (2.11) imply (2.5) for the case $\lambda\to\infty$. In the same way we deduce (2.5) for the case $\lambda\to-\infty$.

Relation (2.6) follows from (2.5) and from the equality (see [8])

$$-\mathrm{i}\lambda\widetilde{s_1}(\lambda) = -\mathrm{i}\lambda\widetilde{s}(\lambda)+\mathrm{e}^{\mathrm{i}\lambda\omega}\bigl(-\ln\omega\bigr)^{-\beta}+o(1), \quad \lambda\to\infty. \tag{2.12}$$

The theorem is proved. □

Thus, the function $-\mathrm{i}\lambda\widetilde{s}(\lambda)$ is bounded. Hence, we obtain the following assertion (see [8]).

Corollary 2.2 *If* $\Re\beta\geq 0$ *and* $0<\omega<1$*, then the operator* S_β *defined by formula* (2.2) *is bounded in the space* $L^2(0,\omega)$.

Further we need the following asymptotic relation (see [8]):

$$\bigl\|S_\beta\mathrm{e}^{-\mathrm{i}x\lambda}+\mathrm{i}\lambda\widetilde{s_1}(\lambda)\mathrm{e}^{-\mathrm{i}x\lambda}\bigr\|\to 0, \quad \lambda\to\infty. \tag{2.13}$$

Using (2.6) and (2.13) we derive two statements below.

Corollary 2.3 *If* $\Re\beta=0$ *and* $|z|=1$*, then the points* z *belong to the spectrum of the operator* S_β.

Corollary 2.4 *If* $\Re\beta=0$, $\nu>0$, $\gamma=\beta+\nu$*, then in the space* $L^2(0,\omega)$ *we have*

$$S_\gamma\to S_\beta, \quad \nu\to 0 \quad \text{(strong convergence)}. \tag{2.14}$$

2. Taking into account (1.3) we can reformulate Corollaries 2.2–2.4 for the operator V_β.

Corollary 2.5 *If* $\Re\beta\geq 0$ *and* $0<\omega<1$*, then the operator* V_β *is bounded in the space* $L^2(0,\infty)$*.*

Corollary 2.6 *If* $\Re\beta = 0$ *and* $|z| = 1$*, then the points* z *belong to the spectrum of the operator* V_β*.*

Corollary 2.7 *If* $\Re\beta = 0$*,* $\nu > 0$*,* $\gamma = \beta+\nu$*, then in the space* $L^2(0,\infty)$ *we have*

$$V_\gamma \to V_\beta, \quad \nu\to 0, \quad \text{(strong convergence)}. \tag{2.15}$$

3. Let us introduce the operator

$$T_\beta f = \int_0^x f(t)\big|\ln(x-t)\big|^{-\beta}\,\mathrm{d}t, \quad \beta > 0, \tag{2.16}$$

where $f(x)\in L^2(0,\omega)$ and $0<\omega<1$.

Proposition 2.8 *The operator* T_β *is such that* $T_\beta\in\sigma_2$ *and* $T_\beta\notin\sigma_1$*.*

Proof. It is obvious that $T_\beta\in\sigma_2$. In order to prove that $T_\beta\notin\sigma_1$ we write the operator T_β in the form

$$T_\beta f = \frac{\mathrm{d}}{\mathrm{d}x}\int_0^x f(t)R(x-t)\,\mathrm{d}t, \tag{2.17}$$

where

$$R(x) = \int_0^x \ln^{-\beta}(u)\,\mathrm{d}u. \tag{2.18}$$

Integrating by parts the right-hand side of the expression

$$\widetilde{s}_1(\lambda) = \int_0^\omega \mathrm{e}^{\mathrm{i}t\lambda}R(t)(1-t/\omega)\,\mathrm{d}t, \tag{2.19}$$

we obtain

$$\widetilde{s}_1(\lambda) = \frac{-1}{\mathrm{i}\lambda}\int_0^\omega \mathrm{e}^{\mathrm{i}t\lambda}\big|\ln(t)\big|^{-\beta}\,\mathrm{d}t. \tag{2.20}$$

Comparing relations (2.3) and (2.20), we see that $-\mathrm{i}\lambda\widetilde{s}_1(\lambda)$ for the operator T_β is equal to $\widetilde{s}(\lambda)$ for the operator S_β. Relations (2.5) and (2.20) imply that

$$\widetilde{s}_1(\lambda) = \frac{\mathrm{e}^{\mathrm{i}\omega\lambda}}{\lambda^2}(-\ln\omega)^{-\beta}\big(1+o(1)\big), \quad \lambda\to\infty. \tag{2.21}$$

It follows from (2.13) and (2.21) that

$$\sum_{n=1}^{\infty}\big|(T_\beta\varphi_n,\varphi_n)\big| = \infty, \tag{2.22}$$

where $\varphi_n(x) = \mathrm{e}^{\mathrm{i}xn/\omega}$.

Next, we use the following result [3]: *if an operator A is compact in the Hilbert space and φ_j is an orthonormal system, then*

$$\sum_{1}^{n}\big|(A\varphi_j,\varphi_j)\big| \le \sum_{j=1}^{n} s_n, \tag{2.23}$$

where $s_j(A)$ is the non-decreasing sequence of the eigenvalues of the operator $(AA^*)^{1/2}$. The formulated result implies the following statement: *if the equality*

$$\sum_{1}^{\infty}\big|(A\varphi_j,\varphi_j)\big| = \infty \tag{2.24}$$

is valid, then $\sum_{j=1}^{\infty} s_j = \infty$. In other words, (2.22) yields

$$\sum_{j=1}^{\infty} s_j = \infty. \tag{2.25}$$

Hence, the proposition is proved. □

3 Friedrichs model

Let us consider in the Hilbert space $L^2(0,\omega)$ the operator

$$A_{\alpha,\beta}f = V_\beta Q V_\alpha f, \quad Qf = xf, \quad \Re\alpha > 0, \quad \Re\beta > 0. \tag{3.1}$$

Using relations (1.1) and (1.2) we obtain

$$A_{\alpha,\beta}f = \int_0^x f(u)U_{\alpha,\beta}(x,u)\,\mathrm{d}u, \tag{3.2}$$

where

$$U_{\alpha,\beta}(x,u) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^{x-u} E_\beta(x-u-y)(u+y)E_\alpha(y)\,\mathrm{d}y. \tag{3.3}$$

We need the relation (see [1]):

$$\int_0^x (x-y)^{\alpha-1}y^{\beta-1}\,\mathrm{d}y = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}x^{\alpha+\beta-1}, \ \Re\alpha > 0, \ \Re\beta > 0. \tag{3.4}$$

Relations (1.2) and (3.4) imply that

$$\frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^{x-u} E_\beta(x-u-y)uE_\alpha(y)\,\mathrm{d}y = \frac{1}{\Gamma(\alpha+\beta)}E_{\alpha+\beta}(x-u)u, \tag{3.5}$$

$$\frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^{x-u} E_\beta(x-u-y)yE_\alpha(y)\,\mathrm{d}y = \frac{(x-u)\alpha}{\Gamma(\alpha+\beta+1)}E_{\alpha+\beta}(x-u). \tag{3.6}$$

Relations (3.3) and (3.5), (3.6) imply the following proposition.

Proposition 3.1 *The operator $A_{\alpha,\beta}$ is defined by* (3.2)*, where*

$$U_{\alpha,\beta}(x,u) = \frac{1}{\Gamma(\alpha+\beta)}E_{\alpha+\beta}(x-u)u + \frac{(x-u)\alpha}{\Gamma(\alpha+\beta+1)}E_{\alpha+\beta}(x-u). \tag{3.7}$$

Using again relations (1.2) and (3.4) we have

$$\frac{1}{\Gamma(m)}V_\beta E_m(y) = \frac{1}{\Gamma(m+\beta)}E_{m+\beta}(x), \quad \Re\beta > 0, \quad \Re m > 0. \tag{3.8}$$

Hence, the equality

$$\lim_{\beta\to 0} V_\beta\left(\frac{1}{\Gamma(m)}E_m(y)\right) = \frac{1}{\Gamma(m)}E_m(x) \tag{3.9}$$

is valid.

Remark 3.2 *It is easy to see, that the functions $E_m(x)$ form a complete system in the Hilbert space $L^2(0,\omega)$,*

Thus, we proved the following statement.

Proposition 3.3 *The operator V_β strongly converges to the identity operator I when $\beta \to +0$.*

In view of (3.1), (3.7) and Proposition 3.3 the following theorem is valid.

Theorem 3.4 *Let* $\Re\alpha = 0$, $\alpha{\neq}0$, $\beta = -\alpha$. *Then the operator*

$$A_\alpha = V_\alpha^{-1} Q V_\alpha \tag{3.10}$$

has the form

$$A_\alpha f = x f(x) + \alpha \int_0^x (x-y) E_0(x-y) f(y)\,\mathrm{d}y, \quad f(x){\in}L^2(0,\omega), \tag{3.11}$$

where

$$E_0(x) = \int_0^\infty \frac{1}{\Gamma(s)} \mathrm{e}^{-Cs} s^{-1} x^{s-1}\,\mathrm{d}s. \tag{3.12}$$

It follows from (3.12) that

$$x E_0(x) = \frac{1}{|\ln x|}\big(1 + o(1)\big), \quad x{\to}0. \tag{3.13}$$

Let us formulate the analogue of the Theorem 3.4 ([9, Ch. 3, Section 4]).

Theorem 3.5 *Let* $\Re\alpha = 0$, $\alpha{\neq}0$. *Then the operator*

$$B_\alpha = J_\alpha^{-1} Q J_\alpha \tag{3.14}$$

has the form

$$B_\alpha f = x f(x) + \alpha \int_0^x f(y)\,\mathrm{d}y, \quad f(x){\in}L^2(0,\omega). \tag{3.15}$$

We note, that the operators A_α and B_α are partial cases of the Friedrichs model [2].

4 Generalized wave operators

1. Let us introduce the notion of the generalized wave operators ([7]).

Definition 4.1 *Let the operators* A *and* A_0 *act in the Hilbert space* H, *where the operator* A_0 *is self-adjoint with absolutely continuous spectrum. We assume that there exists a unitary operator function* $W_0(t)$ *satisfying the following conditions:*

1. The limits in the sense of strong convergence

$$W_{\pm}(A, A_0) = \lim_{t \to \pm\infty} \big(\mathrm{e}^{iAt} \mathrm{e}^{-iA_0 t} W_0(t) \big) \tag{4.1}$$

exist.

2.

$$\lim_{t \to \pm\infty} W_0^{-1}(t + \tau) W_0(t) = I. \tag{4.2}$$

3. The commutations relations hold for arbitrary values t and τ:

$$W_0(t) A_0 = A_0 W_0(t), \quad W_0(t) W_0(t + \tau) = W_0(t + \tau) W_0(t). \tag{4.3}$$

The operators $W_{\pm}(A, A_0)$ are named generalized wave operators.
If $W_0(t) = I$ then the operators $W_{\pm}(A, A_0)$ are usual wave operators.

The formulated notions of wave operators and generalized wave operators are correct and useful not only for self-adjoint operators A, but for non-self-adjoint operators A too.

Definition 4.2 *The spectrum of non-self-adjoint operator A is absolutely continuous if the operator A can be represented in the form*

$$A = V^{-1} A_0 V, \tag{4.4}$$

where the operators V and V^{-1} are bounded and the operator A_0 is self-adjoint with absolutely continuous spectrum.

Proposition 4.3 *Let conditions* (4.1)–(4.3) *be fulfilled. Then*

$$W_{\pm}(A, A_0) \mathrm{e}^{\mathrm{i} A_0 t} = \mathrm{e}^{\mathrm{i} A t} W_{\pm}(A, A_0) \tag{4.5}$$

Proof. As in the of self-adjoint case we use the relation

$$W_{\pm}(A, A_0) = \lim_{t \to \pm\infty} \big(\mathrm{e}^{\mathrm{i} A (t+s)} \mathrm{e}^{-\mathrm{i} A_0 (t+s)} W_0(t + s) \big). \tag{4.6}$$

The second relation of (4.3) and (4.6) imply

$$W_{\pm}(A, A_0) = \mathrm{e}^{\mathrm{i} A s} W_{\pm}(A, A_0) \mathrm{e}^{-\mathrm{i} A_0 s}. \tag{4.7}$$

This proves the proposition. □

Example 4.4 *Let us consider the operator* A_α*, where* $\Re\alpha = 0$*,* $\alpha{\neq}0$*, and the operator* $A_0 = Q$ *(see Theorem 3.4).*

According to equality (3.10) the operator A_α has absolutely continuous spectrum. The following statement is valid.

Proposition 4.5 *We assume that*

$$W_0(t) = \big(\ln^{-\alpha}|t|\big)I. \tag{4.8}$$

We have

$$W_\pm(A_\alpha, Q) = V_\alpha^{-1}. \tag{4.9}$$

Proof. Relations (4.2) and (4.3) are fulfilled. Now, we write the equality

$$\mathrm{e}^{\mathrm{i}A_\alpha t}\mathrm{e}^{-\mathrm{i}Qt}W_0(t)\mathrm{e}^{\mathrm{i}x\tau} = V_\alpha^{-1}\mathrm{e}^{\mathrm{i}Qt}V_\alpha\mathrm{e}^{-\mathrm{i}Qt}W_0(t)\mathrm{e}^{\mathrm{i}x\tau}. \tag{4.10}$$

Using (2.6), (2.13), (4.2) we obtain (4.9). The proposition is proved. □

Example 4.6 *Let us consider the operator* B_α*, where* $\Re\alpha = 0$*,* $\alpha{\neq}0$*, and the operator* $A_0 = Q$ *(see Theorem 3.5).*

According to equality (3.14) the operator B_α has absolutely continuous spectrum. The following statement is valid.

Proposition 4.7 *Assume that*

$$W_0(t) = \big(|t|^{-\alpha}\big)I. \tag{4.11}$$

Then we have

$$W_\pm(B_\alpha, Q) = \mathrm{e}^{\pm(\mathrm{i}\alpha\pi/2)}J^{-\alpha}. \tag{4.12}$$

Proof. Relations (4.2) and (4.3) are fulfilled. For the operator J^α ($\Re\alpha = 0$, $\alpha{\neq}0$) the following relation

$$-\,\mathrm{i}\lambda\widetilde{s_1}(\lambda) = |\lambda|^{-\alpha}\big(1 + o(1)\big)\mathrm{e}^{\pm(\mathrm{i}\alpha\pi/2)}, \quad \lambda\to\pm\infty \tag{4.13}$$

holds (see [8, formula (22)]). Now, we write the equality

$$\mathrm{e}^{\mathrm{i}B_\alpha t}\mathrm{e}^{-\mathrm{i}Qt}W_0(t)\mathrm{e}^{\mathrm{i}x\tau} = J^{-\alpha}\mathrm{e}^{\mathrm{i}Qt}J^\alpha\mathrm{e}^{-\mathrm{i}Qt}W_0(t)\mathrm{e}^{\mathrm{i}x\tau}. \tag{4.14}$$

Hence, using (2.13), (4.13) and (4.2) we obtain (4.12). The proposition is proved. □

References

[1] H. Bateman and A. Erdelyi, *Higher transcendental functions, I*, New York-London, Mc Graw-Hill Book Company, 1953.

[2] L.D. Faddeev, *On a model of Friedrichs in the theory of perturbations of the continuous spectrum,* Amer. Math. Soc. Transl. (2) **62**, 177–203, 1967.

[3] Ky Fan, *Maximum properties and inequalities for the eigenvalues of completely continuous operators*, Proc. Nat. Acad. Sci. USA **37**:11 (1951), 760–766.

[4] I. Gohberg and M.G. Krein, *Theory and applications of Volterra operators in Hilbert space*, Amer. Math. Soc., Providence, 1970.

[5] E. Hille and R. Phillips, *Functional analysis and semigroups*, Amer. Math. Soc., Providence, 1957.

[6] A.I. Markushevich, *Theory of functions of a complex variable, 3*, AMS Chelsea Publishing, 1977 (translated from Russian).

[7] L.A. Sakhnovich, *Dissipative operators with absolutely continuous spectrum*, Trans. Mosc. Math. Soc. **19** (1968), 233–297.

[8] L.A. Sakhnovich, *Triangular integro-differential operators with difference kernels*, Sib. Math. Journ. **19**:4 (1978), 871–877 (Russian).

[9] L.A. Sakhnovich, *Integral equations with difference kernels on finite intervals,* 1st edition, Operator Theory Adv. Appl., vol. 84, Birkhäuser, Basel–Boston–Berlin, 1996.

[10] L.A. Sakhnovich, *On the notion and important classes of (S+N)-triangular operators*, arXiv:1501.02831.

[11] E.C. Titchmarsh, *Introduction to the theory of Fourier integrals*, Oxford, 1937.